

MANTELLICERAS SAXBII, AND THE HORIZON OF THE MARTIMPREYI ZONE IN THE CENOMANIAN OF ENGLAND

by W. J. KENNEDY and J. M. HANCOCK

ABSTRACT. Synonyms of *Mantelliceras saxbii* (Sharpe) include *M. hyatti* Spath and *M. martimpreyi* (Coquand) *sensu* Pervinquière. In England this species occurs at a recognizable horizon that is in the middle, not at the base of the Lower Cenomanian.

SPECIFIC names for Cenomanian ammonites have often been erected haphazardly, ignoring the possibility that specimens of different size may be no more than different growth stages of the same species. The problem is acute in the Cenomanian stage because the ammonites are usually preserved in two quite distinct facies: (i) in limestones (e.g. South Africa, Utatur division of southern India, Chalk of north-west Europe) they occur as moulds preserving the adult body chamber and the later septate portions; (ii) in clays and marls (e.g. Texas, Algeria, and Tunisia) they occur as pyritic or limonitic internal moulds, only preserving minute nuclei or the adults of genuinely tiny species. In southern England the basement beds of Chalk contain ammonites preserved as phosphatic internal moulds and casts, and this preservation sometimes bridges the gap in growth stages between the pyritic nuclei and limestone body chambers. Hence it is possible to demonstrate the equivalence of *Ammonites saxbii* Sharpe (1857, p. 45, pl. 20, figs. 3*a*, *b*). *Mantelliceras martimpreyi* (Coquand) *sensu* Pervinquière (1910, p. 41, pl. 4, figs. 3–9), *Mantelliceras hyatti* Spath (Sharpe 1857, pl. 18, figs. 4*a*, *b*), and one of the syntypes of *Mantelliceras batheri* Spath (Mantell 1822, pl. 22, fig. 1).

It has been pointed out that 'aff.' in this paper is given a different meaning from the interpretation of some other writers. By prefixing a specific name with 'aff.' we mean that the specimen(s) referred to differs rather clearly in some respect(s) from the holotype (or lectotype or neotype) of the species but is still within the limits of variation of the species. It is a notation which allows one to indicate qualitative variation within the pre-evolutionary convention of Linnean nomenclature. This usage was discussed by Trueman and Weir (1946, pp. xx–xxi). We do not mean by 'aff.' that the specimen is necessarily specifically different.

SYSTEMATIC DESCRIPTION

Family ACANTHOCERATIDAE Hyatt 1900

Genus MANTELLICERAS Hyatt 1903

Mantelliceras saxbii (Sharpe)

Plate 79, figs. 1–5; Plate 80, figs. 1–4; Plate 81, figs. 1, 4, 6, 7, 8; Plate 82, figs. 2, 4, 5

- ? 1814 *Ammonites mantelli* J. Sowerby, p. 119 (pars.), pl. 55, upper figure only.
- 1822 *Ammonites mantelli* var. *costata* Mantell, p. 114 (pars.), pl. 22, fig. 1 only.
- 1857 *Ammonites saxbii* Sharpe, p. 45, pl. 20, figs. 3*a*, *b*.
- 1857 *Ammonites mantelli* Sowerby; Sharpe, p. 40, pl. 18, figs. 4*a*–*b* only (*non* Sowerby).
- 1860 *Ammonites saxbii* Sharpe; Pictet and Campiche, p. 320.

- 1864 *Ammonites saxbii* Sharpe; Stoliczka, p. 85.
 1862 *Ammonites martinpreyi* Coquand, p. 172, pl. 1, figs. 7–8.
 ? 1871 *Ammonites mantelli* Sowerby; Schlüter, p. 12, pl. 6, figs. 1, 2, 11.
 1903 *Mantelliceras couloni* (d'Orbigny); Hyatt, p. 114.
 1907 *Acanthoceras martinpreyi* Coquand; Pervinquièrre, p. 289, pl. 16, figs. ?2–3 only.
 1910 *Acanthoceras martinpreyi* Coquand; Pervinquièrre, p. 41, pl. 4, figs. 3–8 only; fig. 9?
 1925 *Mantelliceras hyatti* Spath, p. 197.
 1926 *Mantelliceras batheri* Spath, p. 431 (pars.).
 ? 1929 *Acanthoceras* (*Mantelliceras*) *martinpreyi* Coquand; Collignon, p. 10, pl. 3, figs. 4–5.
 non 1937 *Mantelliceras hyatti* Spath; Collignon, p. 55, pl. 4, fig. 5, pl. 9, fig. 9 (= ? *Eucalycoceras* sp. nov.)
 1937 *Mantelliceras saxbii* (Sharpe); Spath, p. 279.
 1951 *Mantelliceras batheri* Spath; Wright and Wright, p. 24.
 1951 *Mantelliceras hyatti* Spath; Wright and Wright, p. 24.
 1951 *Mantelliceras saxbii* (Sharpe); Wright and Wright, p. 25.
 1963 *Mantelliceras* (*Mantelliceras*) *hyatti* Spath; Renz, p. 1102, pl. 5, figs. 4–5.
 1963 *Mantelliceras* (*Mantelliceras*) sp. aff. *saxbyi* (Sharpe); Renz, p. 1103, pl. 5, fig. 1.
 non 1963 *Mantelliceras* (*Mantelliceras*) *saxbyi* (Sharpe); Renz, p. 1104, pl. 4, fig. 1.
 ? 1963 *Mantelliceras* (*Mantelliceras*) *batheri* Spath; Renz, p. 1107, pl. 5, fig. 3.
 1964 *Mantelliceras hyatti* Spath; Collignon, p. 71, pl. 340, figs. 1518, ?1520.
 non 1964 *Mantelliceras saxbyi* Sharpe; Collignon, p. 98, pl. 341, fig. 1558.
 ? 1967 *Mantelliceras albanense* Busnardo *et al.*, p. 222, pl. 12, figs. 1–2.
 ? 1967 *Mantelliceras* cf. *hyatti* Spath; Busnardo *et al.*, p. 222.
 ? 1967 *Mantelliceras* (*Couloniceras*) *cressierense* Renz; Busnardo *et al.*, p. 223, pl. 12, figs. 3–4.
 ? 1967 *Mantelliceras* (*Couloniceras*) cf. *cressierense* Renz; Busnardo *et al.*, p. 223.

Lectotype. GSM Geol. Soc. coll. 7763, the original of *Ammonites saxbii* Sharpe 1857, pl. 20, figs. 3a–b, from the Lower Chalk of Ventnor, Isle of Wight; designated by Wright and Wright (1951, p. 38).

This specimen is a small, well-preserved, wholly septate, phosphatic internal mould, retaining traces of matrix which is grey, slightly glauconitic chalk. It is moderately involute, just under half the previous whorl being covered. The intercostal whorl section is compressed, with flat sides and a narrowly rounded venter. The costal section is rather similar, with the greatest breadth at the umbilical bulla. The umbilicus is small and shallow, with a rounded umbilical wall.

There are 28 ribs on the outer whorl. Up to a diameter of 18 mm these ribs are very weak; long, flat

EXPLANATION OF PLATE 79

All figures are of natural size. Specimens are coated with ammonium chloride.


Figs. 1a–b. *Mantelliceras saxbii* (Sharpe) (the holotype of *Mantelliceras hyatti* Spath, from the Chalk Marl (Mantelli Zone) of Bonchurch, Isle of Wight. Side and ventral views of BMNH 50288; figured by Sharpe 1857, pl. 18, figs. 4a–b.

Figs. 2a–b. *Mantelliceras* aff. *saxbii* (Sharpe) (the syntype of *Mantelliceras batheri* Spath figured by Mantell 1822, pl. 22, fig. 1, as *Ammonites mantelli* var. *costata*), from the Lower Chalk of Lewes, Sussex. Side, front, and ventral views of BMNH 5694.

Fig. 3. *Mantelliceras saxbii* (Sharpe); phosphatized internal mould from the phosphate bed (Mantelli Zone, *M. saxbii* assemblage horizon) above the Glauconitic Marl of Gore Cliff, Isle of Wight. Side view of WJK 8761.

Figs. 4a–b. *Mantelliceras* aff. *saxbii* (Sharpe); limonitic internal mould from the Lower Cenomanian 700 m north-east of Koudiat el Assel, 11 km north-north-east of Bou Khadra village, eastern Algeria (9865.2971). Side and front views of authors' collection M6; differs from *M. saxbii* in possessing fewer ribs although the ribbing on the venter is stronger and the tubercles on the shoulders are sharper.

Figs. 5a–b. *Mantelliceras* aff. *saxbii* (Sharpe); phosphatized internal mould from the Glauconitic Marl (Mantelli Zone, *H. carcitanensis* assemblage horizon) at Gore Cliff, Isle of Wight. Side and front views of WJK 9958; more inflated than *M. saxbii*.



KENNEDY and HANCOCK, Cenomanian ammonites

ribs arise at the umbilical shoulder, each with a very faint umbilical bulla, and pass straight up the flank; where the whorl starts to contract the ribs turn gently forwards to meet the shoulder at small, pointed, ventro-lateral tubercles. These long ribs are separated by two or three shorter ribs which arise just below mid-flank, but are otherwise similar to the long ribs.

Beyond 18 mm the ribbing becomes much stronger, each long rib bears a distinct umbilical bulla, small lower ventro-lateral tubercles appear, whilst the upper ventro-lateral tubercles become stronger and clavate as the diameter increases.

In the suture line the first lateral saddle is moderately broad and evenly bipartite; the accessory lobe here is as large as the second lateral lobe (U2). The first lateral lobe is not as long as the external lobe, but has a narrow stem and the end is divided into four narrow and elongate accessory lobes.

Discussion. Sharpe's figure is good, although reversed and somewhat restored.

This species is clearly based on a juvenile specimen. A good idea of the middle growth stages of the species can be gained from the following descriptions of the holotype of *Mantelliceras hyatti* Spath, BMNH 50288 (Pl. 79, figs. 1a–b). This specimen cannot be separated specifically from the lectotype of *M. saxbii* at comparable diameters; it also is a phosphatized, wholly septate, internal mould.

The shell is involute, about half the previous whorl being covered. The whorl section is compressed, the greatest breadth being just above the umbilical bulla. The intercostal section is rounded, and there is a broadly rounded venter in costal section; the whorl sides are slightly inflated.

The umbilicus is small and rather deep, the umbilical wall is steep and undercut, the umbilical shoulder is quite sharply rounded.

There are 36–37 ribs at a diameter of 55 mm. Up to a diameter of 30 mm there are 2 short ribs separating the longer ribs, and such pairs of short ribs sometimes occur up to a diameter of 45 mm. The longer ribs arise at the umbilical seam, with an umbilical bulla, developed to varying degrees on successive ribs. Up to 40 mm diameter the ribs are faintly flexed, swinging forwards across the inner part of the flank, backwards across mid-flank and forwards across the ventro-lateral shoulder, where they broaden slightly. There is an angular change in the costal whorl section a little below the shoulder without lower ventro-lateral tubercles, but strong sharp clavate upper ventro-lateral tubercles are developed on all ribs. The venter is flat with a width about half the whorl thickness, and the upper ventro-lateral tubercles are connected by broad, slightly elevated ribs.

The intercalated short ribs are similar to the long ribs except that they arise at various positions on the sides, from just above the umbilical bulla to the mid-flank, and occur in ones or twos between the long ribs.

Over the last part of the outer whorl, at diameters greater than 45 mm, the long and short ribs alternate regularly.

On some ribs there is a faint indication of a mid-lateral swelling, but never a pronounced tubercle.

A later growth stage of *Mantelliceras saxbii* is illustrated by one of the syntypes of *M. batheri* Spath (BMNH 5694, the original of Mantell 1822, pl. 22, fig. 1, from the Lower Chalk of Sussex, bearing Mantell's original label 'xxii, i') although it is slightly more inflated than the type of *M. hyatti*.

This specimen is a well-preserved, slightly worn and distorted, wholly septate, composite internal mould ('normal' chalk preservation of Jefferies 1962) in hard limestone.

The shell is moderately evolute, about half the previous whorl being covered. The whorl section is compressed, and the greatest breadth is at the umbilical bullae. The intercostal section is rounded; the costal section is also approximately rounded, but there is an angular change on the upper part of the sides and a flattened, broadly convex venter. The umbilicus is narrow and deep, the umbilical wall is overhanging and the umbilical shoulder rounded.

There are 37 ribs at a diameter of 92 mm, alternately long and short; up to a diameter of 50 mm there are sometimes 2 short ribs together. The long ribs arise at the umbilical seam, and develop strong umbilical bullae immediately above the umbilical shoulder. These ribs are flat and rounded, the same width as, or slightly narrower than the interspaces. The ribs pass straight across the sides, developing clavate ventro-lateral tubercles on each side of the narrow venter. The ribs are connected across the venter by a low rounded rib. The shorter, intercalated ribs arise about a third of the way up the sides, and are similar to the long ribs.

The adult body-chamber of *M. saxbii* appears at about 100 mm diameter in British material, and extends for over half a whorl. A crushed chalk example is figured as Pl. 80, figs. 1a–b; a phosphatic

example (slightly more inflated than the type would be, and better referred to as *M. aff. saxbii*) is figured as Plate 81, fig. 1. The ribs are broad and rounded, and one long rib alternates with one (rarely two) shorter ribs. The venter is flattened between distinct, rounded and faintly clavate, ventro-lateral tubercles. Close to the aperture (traces are preserved in the specimen shown in Pl. 80, fig. 2), the ribs weaken and approximate; there are many coarse striae which become distinctly inclined forwards. The aperture itself is simple.

TABLE 1. Measurements in mm of *Mantelliceras saxbii* (Sharpe). The last three specimens are from the phosphate bed above the Glauconitic Marl (Mantelli Zone, *M. saxbii* faunal horizon), Gore Cliff, Isle of Wight.

Collection key: GSM = Geological Survey Museum, Institute of Geological Sciences, London; BMNH = British Museum (Natural History); WJK and JMH = authors' collection.

Measurements of diameter, whorl height, and whorl width have been made between ribs.

Specimen	Diameter	Whorl height	Whorl width	Width of umbilicus	No. of ribs on last whorl	No. of primaries	No. of secondaries
Lectotype							
GSM Geol. Soc. coll. 7763	25.9	11.7	8.6	5.4	28	8 a.	20 a.
<i>M. 'hyatti'</i>							
BMNH 50288	56.0	26.4	19.4	11.6	36 a.	14	22 a.
<i>M. 'batheri'</i>							
BMNH 5694	92.4	44.4	37.7 a.	19.8	36	17 a.	19
= <i>M. aff. saxbii</i>							
WJK 8761	68.3	28.8	23.7	19.3	37	18	19
JMH CC761	43.8 a.	22.1 a.	14.8 a.	8.4			
WJK 8938	54.0	24.8	20.6	13.0			

Characters of Mantelliceras saxbii. The distinctive characters of *M. saxbii* are: (i) when young, a markedly compressed whorl section with flat parallel sides; (ii) when young, the mid-flank decoration is weak, but there are markedly sharp ribs on the shoulders and sharp upper ventro-lateral tubercles; (iii) in youth there are numerous short ribs, and pairs of short ribs are maintained to middle age; (iv) numerous ribs—some 28 on nuclei, increasing to 35–38 in the middle stages, but decreasing again to approximately

EXPLANATION OF PLATE 80

All figures are of natural size. Specimens are coated with ammonium chloride.

Figs. 1a–b. *Mantelliceras saxbii* (Sharpe); composite chalk mould from the upper part of the phosphate bed (Mantelli Zone, *M. saxbii* assemblage horizon) above the Glauconitic Marl, Gore Cliff, Isle of Wight. Side and ventral views of JMH CC760; note the retention of tubercles on the shoulders although the whorl compression and probably the clavateness of the tubercles have been exaggerated by crushing.

Fig. 2. *Mantelliceras saxbii* (Sharpe); composite chalk mould from the phosphate bed (Mantelli Zone, *M. saxbii* assemblage horizon) 3 m above the Glauconitic Marl, Eastbourne, Sussex. Side view of WJK 2689; an adult showing striae between ribs on the body chamber.

Figs. 3a–d. *Mantelliceras saxbii* (Sharpe); the lectotype from the Lower Chalk (Mantelli Zone) of Ventnor, Isle of Wight. GSM Geol. Soc. coll. 7763 (Institute of Geological Sciences photograph).

Figs. 4a–b. *Mantelliceras saxbii* (Sharpe); limonitic internal mould from the Lower Cenomanian 700 m north-east of Koudiat el Assel, 11 km north-north-east of Bou Khadra village, eastern Algeria (9865.2971). Side and front views of authors' collection M5.

Figs. 5a–b. Copies of Coquand's original figures of *Ammonites martinpreyi* (1862, pl. 1, figs. 7–8).

Figs. 6a–c. *Mantelliceras ventnorense* Diener; the holotype from the Chalk Marl (Mantelli Zone) of Ventnor, Isle of Wight. Ventral, side and front views of GSM Geol. Soc. coll. 7759.



KENNEDY and HANCOCK, Cenomanian ammonites

30 on an adult; (v) short ribs and ventro-lateral tubercles are maintained on to the adult body-chamber.

The ratio of the whorl height: whorl width is 1.36 in both the lectotype and the holotype of *M. 'hyatti'*, and this compression immediately distinguishes the species from *M. mantelli* (J. Sowerby). Only occasionally does one get more compressed specimens with a ratio as high as 1.5, but less-compressed specimens ranging away from *M. saxbii* s.s. are common (e.g. Plate 81, figs. 1a–b; ratio 1.17–1.18). Up to diameters of about 25 mm, as in the lectotype itself, the width of the venter is around 0.5 of the whorl width; with further growth the whorl section widens faster than the venter, and the venter width may be as low as 0.34 of the whorl width. In *M. mantelli* the width of the venter is more constant at almost exactly half the whorl width.

THE STATUS OF *MANTELLICERAS MARTIMPREYI*

The most important synonym of *Mantelliceras saxbii* is *M. martimpreyi* (= *Ammonites martimpreyi* Coquand 1862, pp. 172–173, pl. 1, figs. 7–8) because it has been adopted by many authors as the zonal ammonite for the lowest part of the Cenomanian.

The original description and figures of this species are in a rather obscure publication; they are therefore reproduced here as Pl. 80, figs. 5a–b. Coquand's original description is as follows:

15. *Ammonites Martimpreyi* H. Coq. pl. 1, fig. 7 et 8.

Diamètre 21 millimètres.

Coquille comprimée, ornée, en travers et par tours, de 28 à 30 petites côtes inégales, dont les unes un peu plus saillantes que les autres, et au nombre de 6 à 8, partent de l'ombilic et viennent se terminer à la périphérie: chacune de ces côtes renferme dans l'intervalle qu'elles laissent entre elles, un nouveau système de côtes, au nombre de 3 à 5, moins nettement accusées et qui vont en s'atténuant vers l'ombilic, où elles se montrent bijugées; chaque côte porte vers son extrémité supérieure deux tubercules, dont l'un, plus saillant, est limité franchement vers le dos, et dont l'autre est placé à deux millimètres en dessous. Dos formé par une carène, s'élevant sensiblement au-dessus des tubercules terminaux; ombilic peu ouvert.

Cette espèce, par ses côtes inégales, par les deux tubercules dont elles sont ornées, ainsi que par sa carène, se distingue des autres *Ammonites* du terrain crétacé.

Elle a été découverte par M. Ville, à Bérrouaghia, province d'Alger, dans l'étage rhotomagien. Je me suis fait un devoir de la dédier au général de Martimprey, sous-gouverneur de l'Algérie, qui m'a fourni les plus grandes facilités pour pénétrer dans les parties les moins connues des possessions françaises.

In the summer of 1965, one of us (W. J. K.) was able to examine Coquand's collection in the Geological Institute, Nepstadion Korut, Budapest (part of Coquand's collection is in the National Museum, but none of that material is relevant here).

The specimens are uncatalogued, and are stored by locality. There are nine specimens in one tray bearing the following label:

'*Ammonites martimpreyi* Coquand.

Roth'. [indecipherable symbol] 'Berrouaghia' [indecipherable word] 'Aumaule'.

This presents an immediate problem, for Coquand does not mention more than one specimen in his description, and figures only one, which would thus be the holotype by monotypy. Pervinquière figured seven of these nine ammonites in 1910 and considered one of them (Pervinquière 1910, pl. 4, figs. 7a, 7b; Pl. 81, figs. 7a–b herein) as 'trés

probablement le type figuré dans "Geol. Pal. S. Constantine", pl. 1, fig. 7-8'. Pervinquièr's figure shows this ammonite to have a diameter of about 21 mm (allowing for the over-pyritization visible) which would agree with Coquand's measurement, but Pervinquièr failed to state that his figures are appreciably reduced: the original is larger; moreover, it does not agree with Coquand's description of the decoration.

The main difficulty is that Coquand stated, indeed emphasized, that his specimen had a keel, and this is clearly shown in his figure. If this was an original feature of the shell, *Ammonites martinpreyi* could hardly be a *Mantelliceras*, but it could be referred to *Cottreautes*, a genus with close affinities to *Mantelliceras* (see Sornay 1955). It is true that the smoothly keeled species of *Cottreautes* that have been described from north Africa only possess a mere trace of a keel (e.g. *C. subboulei* Sornay) but the Malagasy species *C. boulei* Collignon is more distinctly keeled. There are also ammonites with a similar *Mantelliceras*-type decoration on the sides but possessing a tuberculated keel which have been referred to *Acompsoceras*, e.g. *A. dubourdieu* Sornay. But the keel may not have been genuine: in our experience limonitic ammonites from clays in north Africa have quite often developed a 'keel' from being crushed, and, as discussed below, several of the 'co-types' of *A. martinpreyi* have such false keels.

Suture lines cannot help because Coquand did not figure one.

This matter can probably never be fully resolved without neotype designation. We therefore select the original in the Coquand collection figured by Pervinquièr 1910, pl. 4, figs. 7a-b (reproduced here as Pl. 81, figs. 7a-b) as neotype of *Ammonites martinpreyi* Coquand.

Of the seven surviving figured specimens, the originals of figs. 4, 6, 7, and 8 are all specifically inseparable from *M. saxbii*, although there is some variation amongst them for instance in the stage at which ribs become prominent across the venter, and the stage where the lower ventro-lateral tubercle begins to weaken. The original of fig. 8 has faint mid-lateral swellings on the long ribs, although no true tubercle is developed; crushing has produced false keels on the originals of figs. 4 and 8.

The neotype (fig. 7) is a perfectly preserved, wholly septate, limonitic internal mould. It is almost the same size as the lectotype of *M. saxbii*, but loses the lower ventro-lateral tubercles earlier, although the costal whorl section continues to be markedly angular at this level on the side. There are up to three short ribs between any two full length ribs.

The original of fig. 9 is very close to *M. saxbii* but differs in having stronger ribbing on the sides (and the venter?), in having a higher ratio of long to short ribs, develops lower ventro-lateral tubercles earlier (and possibly retains them longer), and possesses faint mid-lateral swellings on some of the long ribs.


EXPLANATION OF PLATE 81

All figures are of natural size. Specimens in figs. 1, 11-13 are coated with ammonium chloride.

Figs. 1a-b. *Mantelliceras* aff. *saxbii* (Sharpe); uncrushed and partly phosphatized specimen from the Glauconitic Marl/Popple Bed junction (Mantelli Zone), Dead Maid Pit, Mere, Wiltshire. Side and ventral views of WJK 5341.

Figs. 2-10. Copies of Pervinquièr 1910 pl. 4, figs. 2-10, the 'cotypes' of *Mantelliceras martinpreyi* (Coquand). Pervinquièr's original figure numbers are retained; figs. 7a-b are the neotype; figs. 10a-b are a specimen of *Mantelliceras* gr. *tuberculatum* (Mantell); fig. 4 is generically indeterminate. Originals in the Geological Institute, Nepstadion Korut, Budapest.

Figs. 11a-13b. *Stoliczkaia* (*Stoliczkaia*) spp. juv.; phosphatic specimens from the Dispar Zone ammonite bed, Upper Greensand, south Dorset. C. W. Wright collection, nos. 7450 (11a-c), 7145 (12a-b), 231 38 (13a-b). These specimens show the similarities of nuclei of *S. (Stoliczkaia)* and *M. saxbii*.



KENNEDY and HANCOCK, Cenomanian ammonites

The original of fig. 2 has a genuine notched keel and may be a *Cottreautes*. The original of fig. 10 is better judged from Pervinqui re's figures and, as Pervinqui re noted, can be compared with *M. mantelli*.

The other three specimens preserved in Budapest do not correspond with figs. 3 or 5, and were thus not figured by Pervinqui re.

TABLE 2. Measurements in millimetres of the syntypes of *Ammonites martimpreyi* Coquand in the Geological Institute, Budapest, figured by Pervinqui re 1910, pl. 4; reproduced here in Plate 81, figs. 2-10.

Fig. No.	Identification	Diameter	Whorl height	Whorl width	Width of umbilicus	No. of ribs
2	<i>Cottreautes</i> ?	10	4.5	3		
3	original not found in museum					
4	<i>M. saxbii</i>	20	11	7		
5	original not found in museum					
6	<i>M. saxbii</i>	22	11	8	4	
7	<i>M. saxbii</i> (neotype of <i>A. martimpreyi</i>)	25	12	11	5	34-35
8	<i>M. saxbii</i> crushed and distorted					
9	<i>M. aff. saxbii</i>	36	19	11	7	34-35
10	now in decomposing fragments					

There are many ammonites figured as *Mantelliceras martimpreyi* but not all of them are synonymous with *M. saxbii*. Pervinqui re (1907) discussed *M. martimpreyi* at length and with discernment; he stressed the similarity with Sharpe's var. B of *Ammonites mantelli* which from Sharpe's description would appear to be represented by Sharpe 1856, pl. 18, fig. 4, i.e. the holotype of *M. hyatti*. Pervinqui re also explains correctly the differences from *M. mantelli* which is more inflated and has mid-lateral tubercles. However, Pervinqui re at this time figured a variety of species under the name *Acanthoceras martimpreyi* (1907, pl. 16).

Fig. 1 is closer to *Mantelliceras ventnorense* Diener because the ribbing is flexuous.

Figs. 2 and 3 are juveniles, probably not specifically determinable, but belong to the *M. saxbii* group.

Figs. 4 and 5 are more inflated and probably belong to another species.

Fig. 16 was renamed *Eucalycoceras lymense* by Spath (1926). We have examined the original in the  cole des Mines in Paris and it is a true *Mantelliceras*, but too inflated for *M. saxbii*.

Fig. 17 is difficult to determine.

Fig. 18 was made a syntype of *Mantelliceras batheri* by Spath (1926). We have examined the original in the  cole des Mines in Paris: it is close to *M. costatum* (Mantell) and only differs from Mantell's species in having abrupt beginnings to the long ribs on the umbilical edge, and a sulcate venter at the tuberculate stage.

Proper assessment of all the small pyritic and limonitic specimens from Madagascar, North Africa and Texas must await detailed study of the plentiful material that we have collected stratigraphically from Algeria and Tunisia. As far as can be judged at present not a single limonitic or pyritic ammonite figured by Collignon (1929, 1964) from Madagascar belongs exactly to *M. saxbii*; this is possibly only a reflection of the variability of nuclei, and need not mean that the species is absent from Madagascar. 'Submantelliceras' *worthense* (Adkins) as figured by Adkins (1928, pl. 20, fig. 11) appears inseparable from *M. saxbii* at comparable diameters; Scott (1926) actually called some of Adkins' specimens *Acanthoceras martimpreyi*.

SPECIES COMPARABLE TO *MANTELLICERAS SAXBII**Mantelliceras ventnorense* Diener

This is the most puzzling relative of *Mantelliceras saxbii*. The holotype, by the original designation of Diener (1925, p. 170) is the original of *Anmonites feraudianus* Sharpe (1857, p. 51; pl. 23, figs. 6a–c) *non* d'Orbigny, GSM 7759, from the 'Grey Chalk at Ventnor' (figured here as Pl. 80, fig. 6). This specimen is a small composite internal mould in hard chalk with a limonitic coat. It is distorted into an ellipse (major diameter 28.7 mm), and the venter is abraded over the last third of a whorl. This poor preservation makes comparison with *M. saxbii* difficult, but the decoration is similar. The chief distinctions of *M. ventnorense* appear to be: (i) the much earlier loss of a distinct lower ventro-lateral tubercle (there is the merest trace of such a tubercle at the smallest diameter visible—12 mm?); (ii) the much more flexuous ribbing. Both these features could be exaggerated in the type by the abrasion, crushing and distortion; the apparently greater inflation is almost certainly artificial. However, examples of *Mantelliceras* which are compressed and have flexuous ribbing are known (e.g. Pervinqui re 1907, pl. 16, fig. 1a–b); several authors have described what are probably larger specimens of this species (e.g. *M. ventnorense* in Fabre 1940, p. 236, pl. 8, fig. 8; and *M. ventnorense* in Collignon 1964, p. 87, pl. 347, fig. 1547). We leave *M. ventnorense* here as a separate species, but further work may well indicate that it would be better treated as a subspecies (or variety) of *M. saxbii*.

Mantelliceras couloni (d'Orbigny)

A neotype for this species was designated by Collignon (1937, p. 56, pl. 11) who distinguished it from *M. hyatti* (i.e. *M. saxbii*) in that: (i) the ribs do not reach the centre of the venter so that between the high ventro-lateral tubercles there is a sinus; (ii) the narrowness of the venter combined with the gentle convexity of the flanks produces a perfectly oval intercostal whorl section.

One of us (J. M. H.) has re-examined the neotype (from Lamnay, Sarthe, France), several topotypes and a number of other examples from the Sarthe. They show that up to moderate diameters (50 mm, possibly more) *M. couloni* is not easily distinguished

EXPLANATION OF PLATE 82

All figures are of natural size. Specimens are coated with ammonium chloride.

Figs. 1a–b. *Mantelliceras* aff. *couloni* (d'Orbigny); internal mould with traces of cast of shell from the basement bed of the Wilmington Sands (Mantelli Zone), Hutchins' Pit, Wilmington, south Devon. Ventral and side views of JMH CC337.

Fig. 2. *Mantelliceras saxbii* (Sharpe). Front view of specimen figured in Plate 79, fig. 3.


Figs. 3a–b. *Utaturiceras vicinale* (Stoliczka); internal mould from the Utatur Group, southern India. Side and front views of Oxford University Museum KY 311.

Fig. 4. *Mantelliceras* aff. *saxbii* (Sharpe); phosphatized internal mould from the phosphate bed (Mantelli Zone, *M. saxbii* assemblage horizon) above the Glauconitic Marl, Gore Cliff, Isle of Wight. Front view of WJK 8763; the nucleus is close to the lectotype but the whorl section in the middle stage is appreciably more inflated.

Fig. 5. *Mantelliceras* aff. *saxbii* (Sharpe); phosphatized internal mould with traces of phosphatized shell from the Popple Bed (Mantelli Zone), Dead Maid Pit, Mere, Wiltshire. Ventral view of JMH CC739; angular shoulders with strong tubercles have been retained to a middle growth stage.



1a



1b



2



4



3a



3b



5

from *M. saxbii*: in *M. couloni* there is a narrow venter in relation to the inter-costal whorl thickness (1:4), and umbilical bullae are absent. Around diameters of 70 mm *M. couloni* develops exaggerated ventro-lateral tubercles whilst retaining a narrow venter; the tubercles are only slightly clavate. In some individuals these great tubercles are continued to diameters of 135 mm; in others the adult body chamber begins around 100–110 mm diameter, on which the ventro-lateral tuberculation rapidly disappears and thereafter the venter merely shows a slight flattening.

Not only are the younger stages possibly indistinguishable, but there are all gradations between *M. couloni* and *M. saxbii*. However, the two extremes have not yet been found together at the same horizon, and one is probably the descendant of the other. The exact horizon of the neotype is not recorded, but its lithology combined with the locality of Lamnay suggests that it was from the top third of the Lower Cenomanian.

Wright and Wright (1951) recorded *M. couloni* from south-west England, correctly noting that the English examples are less compressed than the type. Moreover, the tuberculation is probably never so strong. We figure an example (Pl. 82, figs. 1a–b) from Wilmington which is less compressed than the types of both *M. couloni* and *M. saxbii*, but in other features is intermediate between the two species. This specimen has a venter-width 0.36 of the whorl-width.

Utaturiceras vicinale (Stoliczka)

This ammonite has had a somewhat chequered history. Spath (1926) used '*Acanthoceras*' *vicinale* as a zonal index for the top of the Cenomanian, as did Collignon (1959) and Wright (1957). In 1956 Wright made it the type species of a new genus—*Utaturiceras*. As Casey (1960) notes, this Upper Cenomanian age is erroneous (it probably originated with Kossmat's inference of horizon from the matrix), and it is a Lower Cenomanian genus. He concluded that *Utaturiceras* was a synonym of *Mantelliceras*, and *Ammonites vicinalis* could be matched with *Mantelliceras* of the group of *M. saxbii*.

We had been inclined to follow the view of Casey until the revision of *Utaturiceras vicinale* by Matsumoto *et al.* (1966) combined with the features seen in a specimen in the Oxford University Museum (Pl. 82, figs. 3a–b) showed that the species are slightly different. In particular, juveniles of *U. vicinale* have more flexuous ribbing, whilst the middle and adult stages are more compressed, high whorled and with more numerous subdued ribs (40:35) which are very weak across the venter, whilst the width of the venter is less than a third of the inter-costal whorl thickness. These subtle but important differences give *Ammonites vicinalis* a quite distinct appearance. We are still uncertain whether it merits generic or even subgeneric separation from *Mantelliceras*; it could still be an extreme variant of the *M. saxbii* group. Matsumoto and Sarkar were unfortunate in being dependent on specimens from the Isle of Wight for suture lines of *Mantelliceras*. The suture line differences they list are not so distinct when one compares *U. vicinale* with well preserved *Mantelliceras* limestone internal moulds from Devon (text-fig. 1). For these reasons we are sorry that Matsumoto (1969, p. 291) should have felt it necessary to place *Utaturiceras* in a separate subfamily (Utaturiceratinae) from *Mantelliceras*.

Other species described from Europe

Mantelliceras cressierense Renz (1963, p. 1105, pl. 4, figs. 3a, 3b), from the Swiss Jura, is based on a single crushed specimen; it differs from *M. saxbii* in its coarser, fewer ribs

(29:37 per whorl at 70 mm diameter), but is probably a closely allied species. Renz's *M. saxbii* (pl. 4, fig. 1) differs from that species by its possession of a marked mid-lateral tubercle, but his *M. aff. saxbii* and *M. hyatti* are both *M. saxbii*, as is possibly his *M. ventnorense* (pl. 4, fig. 2).

Busnardo (in Busnardo *et al.* 1966) has figured several *Mantelliceras* from the French Jura whose preservation is rather poor and which are therefore difficult to assess properly. His *M. albanense* (pl. 12, figs. 1, 2) resembles the adult of the more inflated *M. saxbii* such as we figure as Plate 81, fig. 1, but the inner whorls of *M. albanense* are said to have ribs of equal length.

TEXT-FIG. 1. External suture-lines above the umbilical shoulder of *Mantelliceras* and *Utaturiceras*. All $\times 2$. Figs. *b*, *d-f* drawn by J. M. H.

a. Mantelliceras saxbii (Sharpe) Kyushu University no. 9307, T. Matsumoto collection from the Glauconitic Marl, St. Catherine's Point, Isle of Wight. After Matsumoto in Matsumoto, Sastry, and Sarkar 1966.

b. Utaturiceras vicinale (Stoliczka) Oxford University Museum KY 311 from the Utatur Group of southern India.

c. Utaturiceras vicinale (Stoliczka) Lectotype, Geological Survey of India 190, from the Utatur Group, Odium, southern India. After Matsumoto in Matsumoto, Sastry, and Sarkar 1966.

d. Mantelliceras aff. saxbii (Sharpe) JMH CC461 from the *Mantelliceras*-rich bed, Mr. Hutchins' pit, Wilmington, Devon.

e. Mantelliceras saxbii (Sharpe) WJK 8761 from the upper phosphate bed, Gore Cliff, Isle of Wight.

f. Mantelliceras ventnorense Diener JMH CC459 from the *Mantelliceras*-rich bed, Mr. Hutchins' pit, Wilmington, Devon.

In the list below of the features which Matsumoto and Sarkar (1966) say distinguish the sutures of *Utaturiceras* from those of *Mantelliceras*, the exceptions are noted afterwards in brackets:

- (1) They are fairly deeply incised (also seen in fig. *d*);
- (2) There are numerous descending auxiliaries (also seen in fig. *d*).

In addition they note in the suture lines of *Utaturiceras*:

- (3) The deeply bipartite external saddle (also seen in fig. *e*);
- (4) The first lateral lobe is deeper than broad and is bipartite at the bottom; its median foliole is much lower than and overhung by the adjacent lateral folioles (also seen in fig. *f*);
- (5) The first lateral lobe is deeper than the external lobe E (also seen in fig. *e* but not shown by the *Utaturiceras* in fig. *b*);
- (6) The inner (i.e. dorsal) branch of the external saddle is slightly taller than the outer, and narrowed by deeply incised lobes (also seen in fig. *f*);
- (7) The saddle between the lateral lobe L and the second lateral lobe U2 is higher than that between the external lobe and the lateral lobe (also seen in fig. *d* and fig. *f*, but barely shown by the *Utaturiceras* in fig. *b*);
- (8) The saddle inside the second lateral lobe U2 and the auxiliary saddles are arranged on a descending line (also seen in fig. *d*);
- (9) The second lateral lobe U2 is much smaller than the lateral lobe L (if anything this is less marked in the *Utaturiceras* fig. *b*).

It has also been our experience that: (*a*) the first lateral saddle (= external saddle) in *Mantelliceras* of the *saxbii* group is broader than in the specimen figured by Matsumoto and Sarkar (fig. *a*); (*b*) the outer side of the lateral lobe is lower on the side than the lower ventro-lateral tubercles.